

The cytoskeletal protein ezrin regulates EC proliferation and angiogenesis via TNF- α -induced transcriptional repression of cyclin A

Raj Kishore, ... , David Goukassian, Douglas W. Losordo

J Clin Invest. 2005;115(8):2297-2297. <https://doi.org/10.1172/JCI22849E1>.

Erratum

Original citation: *J. Clin. Invest.* 115:1030–1038 (2005). doi:10.1172/JCI22849 Citation for this erratum: *J. Clin. Invest.* 115:2297 (2005). doi:10.1172/JCI22849E1 The name of 1 of the authors, David Goukassian, was misspelled. The corrected author list is shown above. We regret this error.

Find the latest version:

<https://jci.me/22849E1/pdf>





Erratum

The cytoskeletal protein ezrin regulates EC proliferation and angiogenesis via TNF- α -induced transcriptional repression of cyclin A

Raj Kishore, Gangjian Qin, Corinne Luedemann, Evelyn Bord, Allison Hanley, Marcy Silver, Mary Gavin, David Goukassian, and Douglas W. Losordo

Original citation: *J. Clin. Invest.* **115**:1785–1796 (2005). doi:10.1172/JCI22849.

Citation for this erratum: *J. Clin. Invest.* **115**:2297 (2005). doi:10.1172/JCI22849E1.

The name of 1 of the authors, David Goukassian, was misspelled. The corrected author list is shown above.

We regret this error.

Erratum

Prevention of obesity in mice by antisense oligonucleotide inhibitors of stearoyl-CoA desaturase-1

Guoqiang Jiang, Zhihua Li, Franklin Liu, Kenneth Ellsworth, Qing Dallas-Yang, Margaret Wu, John Ronan, Christine Esau, Cain Murphy, Deborah Szalkowski, Raynald Bergeron, Thomas Doebber, and Bei B. Zhang

Original citation: *J. Clin. Invest.* **115**:1030–1038 (2005). doi:10.1172/JCI200523962.

Citation for this erratum: *J. Clin. Invest.* **115**:2297 (2005). doi:10.1172/JCI200523962E1.

During the preparation of the manuscript, the file conversion process led to the introduction of errors in the Methods section. The correct temperatures and procedures are listed below.

Measurement of desaturation index. Fatty acid methyl esters were identified with an Agilent 6890 gas chromatograph connected to an HP-5 column (30 m \times 0.32 mm \times 0.25 μ M film thickness) connected to a flame ionization detector set at 250°C. The column and the injector temperature were set to 50°C. The column temperature was increased to 150°C at 4°C/min, increased to 200°C at 3°C/min, held at 200°C for 25 minutes, and then increased to 225°C at 25°C/min and held at 225°C for 10 minutes. Under these conditions, the Δ 9-16:1, 16:0, Δ 9-18:1, and 18:0 methyl esters eluted at 14.2, 14.8, 19.4, and 20.2 minutes, respectively. The desaturation index was calculated as the ratios of Δ 9-16:1/16:0 and Δ 9-18:1/18:0.

We regret these errors.